

Product	Label Contents	Dimensions (WxH)
CINTAC® (UK)	3 kg - Booklet	111 x 140 mm

Active	Product
metsulfuron-methyl	Alias SX
	Cleancrop
	Mondial
	Simba SX
metsulfuron-methyl + tribenuron-methyl	Ally Max SX
	Biplay SX
	Traton SX
metsulfuron-methyl + thifensulfuron-methyl	Avro SX
	Chimera SX
	Concert SX
	Finish SX
	Harmony M SX
	Presite SX
	Refine Max SX
	Mozaic SX
	thifensulfuron-methyl + tribenuron-methyl
	Inka SX
	Ratio SX

* CINTAC® at 0.4 kg/ha may be mixed with up to 180 g/ha Chekker.

** CINTAC® at 0.4 kg/ha may be mixed with up to 200 g/ha Sekator OD.

To combat the risk of the development of resistance, these mixtures and sequences should only be used in conjunction with an effective autumn-applied non-ALS herbicide programme. Do not tank-mix or sequence CINTAC® with other ALS-inhibitors, including sulfonylureas, unless specifically permitted according to the label instructions.

CLEANING OF APPLICATION EQUIPMENT

To avoid damage to crops other than wheat, the application equipment must be thoroughly de-contaminated after application. Immediately after application, clean all application equipment with a proprietary sprayer cleaner (e.g. All Clear Extra) according to the product instructions. Take care to not mix chlorinating and ammonia based products since it may result in the release of toxic gases.

Dispose of the washings safely. DO NOT spray onto a sensitive crop or land intended to be cropped with a sensitive crop. For disposal of washings in the UK, follow the 'Plant Protection Products (Sustainable Use) Regulations' from the Health and Safety Executive'.

FOLLOWING CROPS after normal harvest or crop failure

After normal harvest of a winter wheat crop treated with 0.5 kg/ha CINTAC® winter wheat or winter barley may be sown in the same year as application and spring barley, spring wheat or sugar beet may be sown the following spring. Winter oilseed rape can also be sown in the same year after ploughing but only where just 0.4 kg/ha has been applied to a winter wheat crop. In the event of crop failure, only plant winter or spring wheat in the same season as an application of CINTAC® and if it has been used in tank-mixture or sequence with one of the permitted ALS-inhibitors, check the partner label and use the most restrictive recommendation for succeeding crops.

EXTENSION OF AUTHORISATION

This extension of the authorised use provides for the use of CINTAC® in respect of crops and situations other than those included on the product label (above). Neither the efficacy or the phytotoxicity of the product for which this Extension of authorisation has been granted have been assessed and, as such, the user bears the risk in respect of failures concerning its efficacy and phytotoxicity.

Product	Label Contents	Dimensions (WxH)
CINTAC (UK)	3 kg - Booklet	111 x 140 mm

CONDITIONS OF SUPPLY

All goods supplied by the company are of good quality and we believe them to be fit for purpose. However, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during or after application, which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

®CINTAC is a registered Trademark of Life Scientific Ltd.

All other brands listed are Trade Marks of other manufacturers where proprietary rights may exist.



Product	Label Contents	Dimensions (WxH)
CINTAC (UK)	3 kg - Booklet	111 x 140 mm