

Product	Label Contents	Dimensions (WxH)
KIPOTA (UK)	1 L - Booklet	240 x 149 mm

Mixtures with other sulfonyl-urea & ALS-inhibitor herbicides

Only mix the SU herbicides listed above with KIPOTA and use the full dose of 0.125 L/ha + adjuvant. When seeking to control wild oats in sequence with SU herbicides or other ALS inhibitors such as florasulam, allow an interval of at least 7 days where KIPOTA is applied first and at least 14 days if applied after the application of SU herbicides. Mixture with Lexus Millenium¹ or other SU/ALS herbicides not listed above may result in reduced control of wild oats.

Sequences:

KIPOTA can be used in sequence with hormone herbicides. When it is applied first, allow an interval of at least 7 days before application of the hormone herbicide. When it is applied after hormone herbicides, allow an interval of at least 14 days before application of CMPP-p or 2,4-DB and at least 21 days before application of MCPA or 2,4-D.

FOLLOWING CROPS:

Any broad leaved crop may be sown in the event of crop failure. After 3 weeks, any cereal may be sown and there are no restrictions on permitted crops after a normal harvest.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.