



Before using any tank mixture, consult and comply with the recommendations of the partner products. Each product should be added separately to the bulk of the water in the spray tank and thoroughly mixed before adding the next chemical.

Always use constant agitation of the sprayer tank during mixing, transportation and application. Spray immediately.

NOTES:

When tank mixing CINTAC® + Biopower + Chekker, the max rate of CINTAC® must not exceed 400g/ha and the max rate of Chekker must not exceed 180g/ha

When tank mixing CINTAC® + Biopower + Sekator, the max rate of CINTAC® must not exceed 400g/ha and the max rate of Chekker must not exceed 200g/ha

When tank mixing CINTAC® + Biopower + Liberator, Firestarter or Regatta, the max rate of CINTAC® must not exceed 400g/ha and the max rate of Liberator, Firestarter or Regatta must not exceed 0.3l/ha

When tank mixing CINTAC® plus Herold the max rate of CINTAC® must not exceed 400g/ha

When tank mixing CINTAC® plus Klaxon the max rate of CINTAC® must not exceed 400g/ha and Klaxon must not exceed 50g/ha

The following tank mixes have been tested for physical compatibility with CINTAC® at recommended rates of use and will mix in the sprayer tank. No tests have been undertaken on crop safety or product performance, therefore use is at users own risk.

SPRING TANK MIX SHEET WHICH PROVIDES INFORMATION IN ADDITION TO WHAT APPEARS ON THE PRODUCT LABEL FOR CINTAC®

FOR BEST RESULTS CINTAC® + BIOPOWER SHOULD BE USED ALONE. IF TANK MIXING IS NECESSARY, LIFE SCIENTIFIC WILL SUPPORT THE PHYSICAL COMPATIBILITY OF TANK MIXES OF CINTAC® + BIOPOWER WITH ONLY **ONE** OF THE PRODUCTS LISTED.

TANK MIXES WITH CHLOROTHLONIL CONTAINING PRODUCTS HAVE BEEN SHOWN TO REDUCE EFFICACY AND ARE NOT SUPPORTED BY LIFE SCIENTIFIC

Crop and environment	Minimum indicated dose	Maximum indicated dose	Maximum number of treatments	Label time of application
Wheat harvest	0.2 g/ha	0.3 g/ha	1 per crop	Flag leaf stage of the crop

PLEASE NOTE:

The following tank mixes are supported for physical compatibility with CINTAC® at recommended rates of use and will mix in the sprayer tank. No tests have been undertaken on crop safety. Provided that all product recommendations are adhered to CINTAC® may be applied in mixture with the following pesticides:

Biopower

Amistar

Aviator 235 Xpro

Azoxystar

Boogie

Capalo

Ceando

Centaur

Cyflamid

Epoconazole SC formulations

Fandango

Firefly 155

Flexity

Fortress

Helix

Jaunt

Justice

Mobius

Proline 275

Talius

Torch

Variano Xpro

Zephyr

Alias SX

Ally Max SX

Avro SX

Barton WG

Boxer

Buctril

Calibre SX

Chekker – see notes

Chimera SX

Concert SX

Dakota

Eagle

Finish SX

Galaxy

GF-184

Harmony M SX

Headland Sure-Fire Gold

Herold (until 31 March in year of harvest – see notes)

Hiker

Hunter

Inka SX

Klaxon – see notes

Liberator (until 31 March in year of harvest – see notes)

Mecoprop-p

Mozaic SX

Presite SX

Ratio SX

Refine Max SX

Regatta (until 31 March in year of harvest – see notes)

Sekator – see notes

Simba SX

Slalom

Spitfire

Starane Gold

Starane Vantage

Starane XL

Traton SX

Chlormequat

Strate

Upgrade

Tank mix sheet produced 1st
May 2019.

Cintac is a registered tradename of Life Scientific and contains mesosulfuron-methyl iodosulfuron-methyl-sodium.

All other products are those of other manufacturers where proprietary rights may exist.

Use plant protection products safely. Always read the label and product information before use.

For further product information including warning phrases and symbols refer to www.lifescientific.com