



The following tank mixes are supported for physical compatibility with KIPOTA at recommended rates of use and will mix in the sprayer tank.

No tests have been undertaken on crop safety or product performance, therefore use is at user's own risk.

Before using any tank mixture, consult and comply with the recommendations of the partner products. Each product should be added separately to the bulk of the water in the spray tank and thoroughly mixed before adding the next chemical.

Always use constant agitation of the sprayer tank during mixing, transportation and application. Spray immediately.

PLEASE NOTE:

The following tank mixes are supported for physical compatibility with KIPOTA at recommended rates of use and will mix in the sprayer tank. No tests have been undertaken on crop safety. Provided that all product recommendations are adhered to KIPOTA may be applied in mixture with the following pesticides.

For adjuvants, Life Scientific recommends methylated seed oils in preference

Abacus

Adigor

Adjust

Ally Max SX

Ambarac/Caramba

Anthem

Arizona

Azoxystar/Amistar

Aurelia/Proline

Axial + Adigor

Bio Syl

Probe/Biopower

Calibre SX

Chlormequat

Cintac

Colorado

Comet

Defy

Drill

Eagle

Fandango

Felix

Firebird

Harmony M SX

Herold

Jubilee SX

Justice

Esker/Kestrel

Lambdastar/Hallmark Zeon

Firestarter/Liberator

Mavrik

Mirage

Moddus + chlormequat (1/2 rates of each)

Monkey

Niantic/Atlantis + Probe/Biopower

Omaha

Optimus

Optimus + chlormequat (1/2 rates of each)

Orius P

Phase II

Phoenix

Pico Pro

Pico Stomp

Oraso Pro/Prosaro

Quantum SX

Revolt

Sceptre

Soratel

Starane XL

Stomp Aqua

Sumi-Alpha

Sumir/Boxer

Talius

Terpal

Thor

Toil

Trinestar/Moddus

Unix

Zonor/Folicur

Zypar

PLEASE NOTE:

Tank-mixing with hormone herbicides

Avoid the use of hormone containing herbicides in mixture or sequence with KIPOTA. When KIPOTA is applied before hormone herbicides, allow an interval of 7 days between applications. When KIPOTA is applied after hormone herbicides, allow an interval of 14 days before applying mecoprop-P or 2,4-DB, 21 days before applying MCPA or 2,4-D.

Tank mixing with sulfonyl ureas and other ALS inhibitor herbicides

DO NOT mix KIPOTA with products containing carfentrazone-ethyl, nor with Attribut or Monitor. Only mix the SU herbicides listed with KIPOTA and use the full dose of 0.125 L/Ha + adjuvant. When seeking to control wild oats in sequence with SU herbicides or other ALS inhibitors such as florasulam, allow an interval of at least 7 days where KIPOTA is applied first and at least 14 days if applied after the application of SU herbicides. Mixture with SU/ALS herbicides not listed may result in reduced control of wild oats.

TANK MIX SHEET PRODUCED 06/10/2022

AZOXYSTAR, CINTAC, NIANTIC and LAMBDASTAR are registered trademarks of Life Scientific.

KIPOTA contains clodinafop-propargyl + cloquintocet-mexyl. AZOXYSTAR contains azoxystrobin. AMBARAC contains metconazole. SUMIR contains florasulam. LAMBDASTAR contains lambda-cyhalothrin. CINTAC and NIANTIC contain mesosulfuron-methyl and iodosulfuron-methyl-sodium. All other products are those of other manufacturers where proprietary rights may exist.

Use plant protection products safely. Always read the label and product information before use. For further product information including warning phrases and symbols refer to www.lifescientific.com or email infoUK@lifescientific.com