

Product	Label Contents	Dimensions (WxH)
SUDO MOR (GB)	1L - Booklet	250 x 135 mm

2 Winter Barley

Apply SUDO MOR at 0.4 l/ha between leaf sheath erect stage and second node detectable stage (GS30 – 32).

OR

Apply SUDO MOR at 0.6 l/ha between the flag leaf just visible stage and flag leaf fully emerged stage (GS37 – 39).

3 Spring Barley

Apply SUDO MOR at 0.5 l/ha between the leaf sheath erect stage and the second node detectable stage (GS30 – 32).

4 Winter and Spring Oats

Apply SUDO MOR at 0.4 l/ha between the leaf sheath erect stage and the first node detectable stage (GS30 – 31).

5 Rye, Triticale and Durum Wheat

Apply SUDO MOR at 0.4 l/ha between the leaf sheath erect stage and the second node detectable stage (GS30 – 32).

6 Grassland (Seed Crops)

Apply SUDO MOR at 0.8 l/ha between the leaf sheath erect stage and the first node detectable stage (GS30 – 31).

MIXING AND SPRAYING

1 Spray Volume

Apply SUDO MOR in a recommended 200 l/ha of water. An increase in water volume will result in increased penetration – this is dependent on crop stage and habit.

2 Mixing and Spraying

Before spraying it is important to check all hoses, filters and nozzles, and to ensure that the sprayer is clean and correctly set to give an even application at the correct volume. Half fill the spray tank with clean water. Begin agitation and add the required quantity of SUDO MOR directly to the tank. Add the remainder of the water and agitate the mixture thoroughly before and during spraying.

Take particular care to avoid overlapping of spray swaths.

Wash out containers with an integrated pressure rinsing device or manually rinsing three times and add the washings to the spray tank. Do not leave the diluted spray in the tank for extended periods such as meal breaks or overnight.

3 Spray Nozzles

Application should be made using a MEDIUM quality spray as defined by BCPC in a minimum of 200 L/ha. A spray pressure of 2-3 bar is recommended.

Product	Label Contents	Dimensions (WxH)
SUDO MOR (GB)	1L - Booklet	250 x 135 mm

Tank Mixes

SUDO MOR is physically compatible with a range of other products but the efficacy of the mixtures have not been confirmed in trials so use is at the grower's risk. If using tank mixes, unless directed otherwise, the preferred order of addition of products to the tank is: water dispersible granules, wettable powders, suspension concentrates (flowables), emulsifiable concentrates, soluble concentrates. Each product must be added to a half-full sprayer and be fully dispersed before the addition of the next product.

Tank mixes must only be applied within the label recommendations of every product in the mix. Contact your supplier for compatibility information on specific tank mixes. Manufacturer's instructions must be followed for each tank-mix component.

CLEANING OF APPLICATION EQUIPMENT

To avoid damage to other crops, the application equipment must be thoroughly de-contaminated after application.

- Immediately after application, drain the tank completely and wash down with clean water. Rinse out the tank and flush through the booms and hoses.
- Half-fill the tank with clean water and add the recommended dose of detergent cleaner. Agitate and then flush the boom and hoses with the cleaning solution. Top up the tank so that it is completely full and leave to stand for 15 minutes with the agitation running. Flush the booms and hoses again and drain completely.
- Remove the nozzles and filters and clean separately in a solution of detergent cleaner in 10 litres of water.
- Rinse the tank again with clean water, using at least 10% of the tank volume and dispose of the washings safely. For disposal of washings in the UK, follow the 'Code of Practice for Using Plant Protection Products', while in Ireland you should comply with local and national regulations.

CONDITIONS OF SUPPLY

All goods supplied by the company are of good quality and we believe them to be fit for purpose. However, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during or after application, which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

© SUDO MOR is a registered Trademark of Life Scientific Ltd.